

Resistant Starch Content in Foods Commonly Consumed in the United States: A Narrative Review



Mindy A. Patterson, PhD, RDN; Madhura Maiya, PhD; Maria L. Stewart, PhD

ARTICLE INFORMATION

Article history:

Submitted 16 April 2019

Accepted 21 October 2019

Keywords:

Dietary fiber

Food analysis

Resistant starch

Insulin sensitivity

Qualified health claim

2212-2672/Copyright © 2020 by the Academy of Nutrition and Dietetics.

<https://doi.org/10.1016/j.jand.2019.10.019>

ABSTRACT

Resistant starch (RS; types 1 to 5) cannot be digested in the small intestine and thus enters the colon intact, with some types capable of being fermented by gut microbes. As a fiber, types 1, 2, 3, and 5 are found naturally in foods, while types 2, 3, and 4 can be added to foods as a functional ingredient. This narrative review identifies RS content in whole foods commonly consumed in the United States. Scientific databases ($n=3$) were searched by two independent researchers. Ninety-four peer-reviewed articles published between 1982 and September 2018 were selected in which the RS was quantified and the food preparation method before analysis was suitable for consumption. The RS from each food item was adjusted for moisture if the RS value was provided as percent dry weight. Each food item was entered into a database according to food category, where the weighted mean \pm weighted standard deviation was calculated. The range of RS values and overall sample size for each food category were identified. Breads, breakfast cereals, snack foods, bananas and plantains, grains, pasta, rice, legumes, and potatoes contain RS. Foods that have been cooked then chilled have higher RS than cooked foods. Foods with higher amylose concentrations have higher RS than native varieties. The data from this database will serve as a resource for health practitioners to educate and support patients and clients interested in increasing their intake of RS-rich foods and for researchers to formulate dietary interventions with RS foods and examine associated health outcomes.

J Acad Nutr Diet. 2020;120(2):230-244.

DIETARY FIBER CONSISTS OF MANY TYPES OF NON-digestible plant components that have been found to reduce the risk of cardiovascular disease, cancer, and obesity while promoting gut health and immune function.¹ Despite the health-related properties associated with adequate dietary fiber intake, US consumption patterns are substantially lower than the current recommendations of 25 g for females and 38 g for males, or 14 g per 1,000 calories daily.^{2,3} The Institute of Medicine defines dietary fiber as nondigestible carbohydrate and lignin intrinsic and intact in plants, as well as those nondigestible carbohydrates that are isolated and provide a physiological benefit.³ Resistant starch (RS) is a type of fiber that can be categorized as being intrinsic to starchy plants or isolated to function as an ingredient.

RS is the component of starch that is not hydrolyzed by digestive enzymes and enters the large intestine intact, with some types undergoing fermentation by resident microbiota. Because of the digestive properties of RS it is considered an insoluble fiber. Five types of RS have been classified (RS1, RS2, RS3, RS4, and RS5) based on the nature and properties of the RS granule. RS1 includes granules that are resistant, as a result of physical inaccessibility, to digestive enzymes because of the physical nature of the food. RS1 is found in

seeds, legumes, and cereals that are partially milled (eg, whole grains), in which the granule is trapped in the cell wall or matrix. RS2 granules are resistant because of a tightly packed amylose structure that forms a crystallized molecule. RS2 is found in raw potatoes and green bananas, as well as legumes and high-amylose corn. RS2 is resistant to digestion because of its native granule conformation. RS3 is formed when the starch granules are gelatinized and then cooled and causes the starch to crystallize, which is known as retrogradation. RS4 is a chemically modified RS that is not naturally occurring in foods, but can be added as a functional ingredient to improve the dietary fiber content in foods. RS5 is heat-stable and formed when lipids bind to amylose in the starch granule to prevent expansion of the granule, which is necessary for digestive enzyme hydrolysis.⁴

Foods with naturally occurring RS include potatoes, cereals, whole grains, beans, legumes, rice, and bananas—many of which are staple foods in the United States. The RS in these foods can be altered because of several factors, including variety, preparation or cooking method, storage temperature and duration, and serving temperature.⁵ For example, higher amylose to amylopectin ratio within the starch granule can compact under certain conditions to become resistant to hydrolysis. RS concentrations can also be influenced by the

environment, natural selection, and mutations.⁴ These inconsistencies, along with the methodology used to quantify RS in foods, make it challenging to identify the amount of RS in foods.

RS can be measured using the integrated dietary fiber methods utilizing chromatography (AOAC 2011.25, 2009.01), enzymatic-gravimetric methods (AOAC 991.43, 985.29), and a specific RS method (AOAC 2002.02); however, the amount of RS quantified by these methods varies, and poses a challenge for labeling and understanding actual dietary RS intake.⁶ Estimation of RS in foods was initially proposed by Englyst and colleagues,^{7,8} who recognized RS fractions in both plant foods (in vitro) and human ileostomy effluent samples (in vivo). In vitro analysis can be either direct or indirect. The amount of RS quantified is dependent on several factors, such as sample preparation, sample pretreatment, enzymes used, and incubation conditions.⁹ The Englyst method indirectly analyzes RS, where the total RS is estimated to be the difference between total starch and sum of rapidly and slowly digested starch.⁷ Modifications to this indirect method include reduction of sample particle size and change in incubation temperature to 37°C, which established a preliminary basis towards quantification of RS in a food sample and further enabled the estimation of different types of RS.¹⁰

Berry¹¹ was one of the first to modify the indirect procedure proposed by Englyst and colleagues^{7,8} to measure RS directly. Modifications proposed by Berry included omitting the initial heating step to 100°C, changes in α -amylase and pullulanase ratio to starch, and a KOH-dependent solubilization procedure. Subsequent changes to both Berry and Englyst procedures by Goni and colleagues,¹² Muir and O'Dea,¹³ and Akerberg and colleagues¹⁴ aimed to replicate the actions in the human digestive tract. Predominant changes in methodology included sample pretreatment (chewed vs milled vs homogenized), enzyme variations (sample treatment with pepsin, replacing pullulanase with amyloglucosidase), and the pH at incubation (5.0 vs 5.2 vs 6.9).

More recently, McCleary and Monaghan¹⁵ reported that omitting the sample treatment with pepsin, inclusion of amyloglucosidase at an incubation pH of 6.0, and a shaking tube experiment resulted in replicating in vivo conditions. A commercially available kit based on this methodology (AOAC 2002.02) is commonly used to determine RS in foods and is available through Megazyme.¹⁶ This kit accurately measures total RS, which is the sum of RS1, RS2, and RS3; however, it is not a reliable measure of RS4 because of underestimation.^{9,16} There are currently no reported methods to estimate RS5 in foods.

Numerous health benefits of RS have been observed in a variety of populations. The resident gut microbiota ferment primarily RS2 and RS3 in the distal bowel to produce short-chain fatty acids (butyrate, propionate, and acetate), which decreases fecal pH.¹⁷ Other gastrointestinal benefits from RS intake include increased fecal wet weight,¹⁷ mild laxation, defecation ease, and frequency.¹⁸ Because some types of RS can be fermented by gut bacteria, RS may be considered a prebiotic.^{19,20} To classify RS as a prebiotic, more evidence that RS selectively increases the abundance of gut bacteria and provides benefit to the host is needed. Studies have shown that RS2 improves *Ruminococcus bromii* and *Eubacterium rectale*²¹ and increases the *Bifidobacteria* genus in middle-

RESEARCH SNAPSHOT

Research Question: What are the foods consumed in the United States that contain resistant starch (RS) and how much RS do they contain?

Key Findings: In this narrative review, foods containing RS were identified, which include breads, cereals, cakes and muffins, chips and snacks, cookies and crackers, bananas and plantains, grains, noodles and pasta, rice, legumes, potatoes, and hazelnuts. Barley, potatoes, and rice with higher amylose concentrations have more RS than lower amylose varieties, and foods cooked then chilled have higher RS than foods cooked without chilling.

aged and elderly adults, while improving dysbiosis from *Proteobacteria* in the elderly adults.²² Another study found that a 2-week supplementation of 66 g RS2/day in adults with reduced insulin sensitivity increased the ratio of Firmicutes to Bacteroidetes,²³ suggesting that RS2 can alter the gut microbiome in populations with metabolic dysfunction. RS also reduces gastric emptying to promote feelings of satiety²⁴ and lowers energy intake at a subsequent meal.²⁵

Consistent data have shown that RS lowers postprandial glycemia and insulinemia^{26,27}; improves insulin sensitivity²⁸⁻³⁰; and reduces inflammation,³¹ postprandial fat oxidation,³² and both serum and low-density lipoprotein cholesterol.³³ A recent review summarized the body of evidence from animal and human trials, specifically noting that the most consistent benefit was related to postprandial blood glucose management and improvements in insulin sensitivity.³⁴ Additional publications have examined the role of RS in digestive health and cardiometabolic risk factors.^{23,31,35}

Many of the health benefits of RS result from RS consumed as a functional fiber in adequate amounts (≥ 15 g/day)^{26,29} with either acute administration³⁶ or over time (≥ 4 to 12 weeks).^{28,37} However, with appropriate meal planning, this level of RS can be achieved through foods with naturally occurring RS. In fact, RS has been attributed to improved health in populations consuming a non-Western diet, such as an African diet rich in corn meal.³⁸ Many RS-containing foods are rich in fiber and can contribute to the total US dietary fiber intake and benefit overall health. The mean RS intakes in the United States have been estimated to be 4.9 g/day (range=2.8 to 7.9 g/day).³⁹ However, the intake data were calculated from 155 individual foods, where only approximately 57% of the foods reported were analyzed using methods that best mimic human digestion.³⁹ Over the past several years, the number of foods analyzed for RS has expanded considerably as a result of a commercially available RS analysis kit utilizing a method that simulates RS digestion. Therefore, the purpose of this narrative review was to create a database using data from peer-reviewed journal articles that quantified the RS content in foods commonly consumed in the United States. In addition, differences in variety, preparation, storage, and processing of RS content in food will be described. The creation of this database is important because it will allow health practitioners and researchers to identify foods containing RS and examine how consuming these foods contributes to health outcomes. The benefits of

RS related to insulin sensitivity have been reported when adequate amounts (approximately ≥ 15 g/day)²⁹ are consumed from RS administered as a functional ingredient, not foods with naturally occurring RS. Additional research is needed to identify the amount and duration of RS found naturally in foods to produce health outcomes. By utilizing the database, 15 g of RS from food can be achieved in free-living individuals with appropriate meal planning.

METHODS

Three databases (PubMed, Scopus, and Science Direct) were utilized to identify peer-reviewed articles published between 1982 and September 2018 that analyzed the amount of RS in foods commonly consumed in the United States for this narrative review. The search terms *resistant starch*, *English language*, and *food** were used to identify references. References were included if the RS analysis was conducted in a laboratory either inside or outside of the United States, but the foods were commonly consumed in the United States. Excluded terms included *human**, *animal**, and *clinical trial**. Exclusion criteria included the use of RS from flours or starches, isolated raw starch, or uncooked foods unsuitable for consumption (eg, raw potatoes and raw cassava). However, raw bananas and plantains and raw oats were included. References that used specific methods not commonly used by consumers before food consumption (eg, adding an acid, enzymes such as pullanase, extrusion of raw material to create an ingredient, or other process) were also excluded. Autoclaved foods were included because this is a technique used in the canning process. Foods that were cooked or milled incompletely were excluded. Foods cultivated to produce a mutant of the native food as a means to alter the nutritional or RS content were also excluded, except for barley, rice, and potatoes, as they were identified as low- or high-amylose varieties. The references meeting inclusion criteria were hand selected by two independent researchers according to the title and abstract, followed by reviewing the full text. In this review, data were pooled from all of the studies that analyzed RS using both in vivo and in vitro methods. Any references that reported either heating the sample before enzyme hydrolysis or an incubation temperature of 100°C were omitted, as it has been reported that this method estimated only RS3 and not RS1 or RS2.⁹

The references presented the RS data in one of two ways: percent dry weight (g RS/100 g dry weight) or as a whole food with the moisture included (g RS/100 g whole food). Both data types were included in the database. However, because the objective of this review is to provide RS content in whole foods, the percent RS in dry weight was adjusted to include moisture content. Moisture was either measured and provided in the article or identified by the US Department of Agriculture FoodData Central database⁴⁰ from the most similar food item. The pooled data were compiled in an Excel (Microsoft) database, then summarized in the Table according to food category.

Statistical Analysis

The raw data added to the Excel database included percent RS based on dry weight (g RS/100 g dry weight) \pm standard deviation (SD), percent RS in the whole food (g RS/100 g whole food) \pm SD, the number of samples analyzed in each reference, the percent moisture either as provided in the reference or as

identified in the US Department of Agriculture FoodData Central database.⁴⁰ Using the raw data, calculations were employed to adjust for moisture content if percent dry weight was given in the reference, as well as the individual weighted means for each item followed by the overall weighted mean \pm weighted SD.

When only percent RS by dry weight was given, the formula $(1 - \text{moisture})\% \text{ dry weight RS}$ was used to adjust for moisture, which provided RS content in the whole food. Due to some references measuring RS in several samples, the weighted mean was calculated for each food item. The weighted mean was determined by multiplying the number of RS samples analyzed in the reference by the whole food RS content (with moisture). Then the overall weighted mean was calculated by summing the individual weighted means and dividing by the total number of samples from all references in the food category. Next, the weighted SD was calculated for each individual food item by multiplying the sample size by the mean RS content in the whole food minus the overall weighted mean squared. The individual weighted SD was summed and divided by the total sample size minus 1, then the square root of this value was calculated to obtain the overall weighted SD. The overall weighted mean of whole food RS and overall weighted SD for each food category are reported in the Table. In addition, the ranges (minimum to maximum) of weighted means in each food category are listed in the Table. Also, the total number of samples in each food category is provided. Note that some references analyzing RS in more than one sample list the mean without the SD. However, the SD for each food item was not necessary to calculate the overall weighted SD. Some references listed a mean RS value without noting the sample size in which a value of 1 was given in the Table. RS values that were $>200\%$ of the mean were removed from the database to control for potential outliers.

RESULTS

The Table provides the overall weighted mean \pm weighted SD, range (minimum to maximum for each food item), and the total sample size of naturally occurring RS concentrations in the whole food for each food item from 94 articles. All foods requiring percent moisture values from the FoodData Central database were available except baked corn tortillas, where the measured value from a reference was used.⁶³ The number of individual food samples included in each food category include 145 bananas and plantains,^{13,41-46} 446 breads,^{8,10,46-80} 37 breakfast cereals,^{10,14,50,79,81,82} 222 ready-to-eat cereals,^{8,10,12-15,41,45,47,49-52,74,75,79,81,83,84} 11 cakes and muffins,^{51,54,85} 18 chips and snacks,⁴⁷ 50 cookies and crackers,^{47,49-52,85} 103 cooked grains,^{10,50,74,76,79,82,84,86-89} 765 legumes,^{10,15,45,50-52,69,75,76,79,90-109} 98 noodles and pasta,^{10,14,41,47,50,52,74,76,78,81,82,110-112} 6 hazelnut,¹¹³ 311 potatoes,^{10,13,14,41,45,50,52,76,80,81,114-122} and 170 rice.^{5,14,41,50,52,81,84,123-127} The foods with the highest mean RS content included uncooked or raw foods, such as uncooked rolled oats (7.7 g/100 g), uncooked rolled milled oats (6.5 g/100 g), ripe raw plantains (5.1 g/100 g), and unripe raw plantains (5.0 g/100 g). Of the cooked foods, high-amylose potatoes that were cooked then chilled for 2 days (6.4 g/100 g); butter or lima beans (6.4 g/100 g); potato salad (5.2 g/100 g); yellow potatoes that have been cooked, chilled, then reheated (5.1 g/100g); corn tortillas that have been

Table. Naturally occurring resistant starch in whole foods commonly consumed in the United States, extracted from peer-reviewed articles published from 1982 to September 2018

Food	g RS ^a per 100 g Food ^b		
	Weighted mean±weighted SD ^{cd}	Range ^e	n ^f
Bananas and plantains			
Banana, unripe, cooking type, raw ^{41,42}	2.8±0.1	2.3-2.9	42
Banana, ripe, cooking type, raw ^{42,43}	1.8±0.1	1.0-2.0	42
Banana, cooked ^{13,44}	0.0±0.0	0.0-0.3	11
Banana, cooked and cooled ⁴⁴	2.0	NA ^g	NA
Banana, cooked, cooled and reheated ⁴⁴	1.4	NA	NA
Plantain, unripe, raw ^{42,45}	5.0±1.1	4.2-7.0	18
Plantain, ripe, raw ^{42,46}	5.1±13.6	1.7-11.4	15
Plantain, cooked ⁴⁴⁻⁴⁶	2.6±1.3	0.0-3.5	10
Plantain, cooked and cooled ⁴⁴	3.2	NA	NA
Plantain, cooked, cooled and reheated ⁴⁴	1.2	NA	NA
Plantain, cooked, unripe chips ⁴⁶	0.0	NA	3
Bread			
Bagel, plain ⁴⁷	0.7	NA	2
Barley, commercial variety ⁴⁸	1.0±0.3	0.7-1.3	4
Barley, high β -glucan ⁴⁸	0.9±0.2	0.8-1.1	2
Barley, low β -glucan ⁴⁸	0.9±0.2	0.8-1.1	2
Barley, low amylose ⁴⁸	0.4±0.0	0.3-0.4	2
Barley, high amylose ⁴⁸	1.6±0.4	1.3-1.9	2
Breadsticks, white wheat ⁴⁷	0.4	NA	2
Brioche ⁴⁹	1.1	NA	NA
Biscuit, oatmeal ⁵⁰	0.9	NA	NA
Biscuit, spelt ⁵¹	0.5	NA	3
Biscuit, unspecified type, store bought ⁵²	1.1	NA	4
Biscuit, wheat, whole grain ⁵¹	0.5	NA	3
Ciabatta, plain ⁵³	0.5	NA	NA
Corn bread ⁵⁴	0.5	NA	NA
Croissant, store bought ⁵³	0.3	NA	NA
Crumpet ^{47,53}	0.7±0.5	0.1-1.0	3
Croutons, store bought ⁴⁷	1.4	NA	2
English muffin, white, store bought ^{47,53}	0.7±0.5	0.1-1.0	3
Focaccia ⁴⁷	1.2	NA	2
French baguette, white, long and thin ^{49,53}	1.0±0.6	0.4-1.4	3
Gluten-free bread ^{55,56}	0.9±0.3	0.5-1.2	15
Mixed-grain bread ^{47,57-59}	2.6±4.9	1.5-4.4	23
Naan ⁵³	0.2	NA	NA
Oatmeal bread ⁴⁷	1.2	NA	2
Pastry shell, store bought, baked ⁴⁷	0.5±0.1	0.4-0.5	4
Pita bread, wheat, store bought ⁴⁷	0.5	NA	2
Pita bread, white, store bought ^{47,53}	0.5±0.4	0.1-0.7	3

(continued on next page)

Table. Naturally occurring resistant starch in whole foods commonly consumed in the United States, extracted from peer-reviewed articles published from 1982 to September 2018 (*continued*)

Food	g RS ^a per 100 g Food ^b		
	Weighted mean±weighted SD ^{cd}	Range ^e	n ^f
Pizza crust, baked ^{47,60}	0.9±0.9	0.4-2.0	3
Potato dumpling ⁶⁰	0.4	NA	NA
Pumpernickel bread ^{47,53}	0.9±0.5	0.4-0.9	3
Rye bread ^{50,57}	3.0±0.2	2.9-3.2	3
Scones, white with fruit ⁵³	0.1	NA	NA
Sourdough bread, spelt ⁵¹	0.7±0.0	0.6-0.7	6
Sourdough bread, wheat ^{61,62}	3.3±3.1	0.7-5.0	10
Sourdough bread, whole wheat ⁵¹	0.7±0.0	0.7-0.7	6
Spelt bread ^{51,61}	0.8±0.0	0.8-0.8	11
Tortilla, corn, store bought ^{h 47}	0.8	NA	2
Tortilla, corn, freshly baked ^{h 63-71}	2.6±0.6	1.8-4.0	33
Tortilla, corn, stored 4°C, 24 h ^{h 65-69}	3.3±0.8	2.1-4.7	24
Tortilla, corn, stored 4°C, 48 h ^{h 65-69}	3.6±0.7	2.5-5.0	24
Tortilla, corn, stored 4°C, 72 h ^{h 65-69}	3.9±0.6	2.9-5.1	24
Tortilla, corn, stored 4°C, 96 h ^{h 65}	4.5±0.5	4.0-5.1	9
Tortilla, corn, stored 4°C, 7 d ^{h 68}	4.6	NA	18
Tortilla, corn, stored 4°C, 14 d ^{h 68}	4.7	NA	18
Tortilla, corn, stored 25°C ^{h 67}	3.6±0.3	3.2-4.0	18
Tortilla, corn, hard shell ^{47,69}	2.7±1.2	0.9-3.9	5
Tortilla, flour, store bought ^{47,53}	0.2±0.2	0.0-0.4	5
Waffles, multigrain, toasted ⁴⁷	0.5	NA	2
Waffles, plain, toasted ⁴⁷	0.6	NA	2
Wheat-germ bread ⁵³	0.1	NA	NA
White bread, added fiber ⁷²	1.7±0.2	1.4-1.9	12
White bread ^{8,10,46,47,50-53,57-62,72-79}	1.0±0.9	0.0-3.9	85
White bread, baked 120°C to 150°C for 3 to 4 h ⁶¹	1.1±0.0	1.1-1.1	8
White bread baked 120°C to 150°C for 12 to 20 h ⁸⁰	2.2±0.5	1.9-2.5	2
White bread, crusty or toasted ^{53,60}	1.9±2.4	0.2-3.6	2
Whole-wheat bread ^{10,47,50,53,77,79}	1.7±1.2	0.3-2.9	14
Breakfast cereal			
Oats, unknown type, cooked ^{10,50}	1.0±1.2	0.1-1.8	2
Oats, flaked, cooked ⁸¹	0.3	NA	6
Oats, rolled, uncooked ⁸²	7.7	NA	NA
Oats, rolled, milled, and uncooked ⁸²	6.5	NA	NA
Rice, cooked ^{14,79,81}	1.0±0.6	1.2-1.6	18
Semolina, cooked ⁸¹	4.8	NA	6
Wheat, cooked ⁷⁹	0.4	NA	3
Breakfast cereal, ready to eat			
Corn, flaked ^{8,10,12-15,41,45,47,49,51,52,74,75,79,83,84}	4.0±1.8	1.0-6.3	165
Corn, puffed ⁴⁷	1.4	NA	2

(continued on next page)

Table. Naturally occurring resistant starch in whole foods commonly consumed in the United States, extracted from peer-reviewed articles published from 1982 to September 2018 (*continued*)

Food	g RS ^a per 100 g Food ^b		
	Weighted mean±weighted SD ^{cd}	Range ^e	n ^f
Corn, square ⁴⁷	1.3	NA	2
Granola ⁴⁷	0.1	NA	2
Multigrain, flaked ⁴⁷	0.0	NA	2
Multigrain, baked ⁴⁷	0.8	NA	2
Oat, bran ^{47,50}	0.5±0.4	0.3-0.7	3
Oat, flaked ^{14,79}	0.6±0.5	0.3-0.7	9
Oat, square ⁴⁷	0.9±0.3	0.6-1.2	4
Rice, flaked or crisped ^{10,45,47,49,50}	1.4±0.8	0.0-2.5	11
Rice, square ⁴⁷	4.2	NA	2
Wheat bran, flaked ^{47,50}	0.8±0.2	0.7-1.1	3
Wheat, puffed ^{50,81}	1.9±1.9	1.2-6.2	7
Wheat, shredded ^{10,47,50}	1.0±0.7	0.0-1.6	4
Wheat, square ⁴⁷	1.4	NA	2
Wheat, whole, flaked ⁴⁷	1.0	NA	2
Cakes and muffins			
Cake, apple crumble ⁵⁴	0.2	NA	NA
Cake, plain, white ⁸⁵	1.8	NA	NA
Cake, unspecified type ⁵⁴	0.4	NA	NA
Muffins, plain ⁸⁵	1.0	NA	NA
Muffins, spelt ⁵¹	0.4	NA	3
Muffins, unspecified type ⁵⁴	1.0	NA	NA
Muffins, wheat ⁵¹	0.6	NA	3
Chips and snacks (store bought)			
Cheese puffs ⁴⁷	0.2	NA	2
Chips, corn, low fat ⁴⁷	0.7	NA	2
Chips, corn ⁴⁷	0.8	NA	2
Chips, multigrain ⁴⁷	0.9	NA	2
Corn puffs, fried ⁴⁷	0.9	NA	2
Granola bar, oats and honey ⁴⁷	0.2	NA	2
Popcorn cakes ⁴⁷	0.3	NA	2
Pretzels ⁴⁷	1.0	NA	2
Rice cakes ⁴⁷	0.2	NA	2
Cookies and crackers (store bought unless specified)			
Biscuit, tea or sugar ^{47,50,52}	1.1±0.5	0.1-1.4	7
Cone, ice cream ⁴⁷	0.3	NA	2
Cone, sugar, old fashioned ⁴⁷	0.5	NA	2
Cookies, ginger snaps ⁴⁷	0.4	NA	2
Cookies, oatmeal ⁴⁷	0.2	NA	2
Cookies, plain, laboratory prepared ⁸⁵	0.8	NA	NA
Cookies, spelt, laboratory prepared ⁵¹	0.2	NA	3

(continued on next page)

Table. Naturally occurring resistant starch in whole foods commonly consumed in the United States, extracted from peer-reviewed articles published from 1982 to September 2018 (*continued*)

Food	g RS ^a per 100 g Food ^b		
	Weighted mean±weighted SD ^{cd}	Range ^e	n ^f
Cookies, vanilla wafers ⁴⁷	0.2	NA	2
Cookies, wheat, laboratory prepared ⁵¹	0.3	NA	3
Crackers, club ⁴⁷	0.4	NA	2
Crackers, graham ⁴⁷	0.3	NA	2
Crackers, multigrain ⁴⁷	1.0±0.6	0.5-1.5	4
Crackers, melba type ⁴⁷	1.2±0.3	0.9-1.5	4
Crackers, rice crunch ⁴⁷	0.4	NA	2
Crackers, rye crisp (store bought and laboratory prepared) ^{47,50}	2.0±2.0	0.9-4.3	3
Crackers, saltine type ⁴⁷	0.6±0.1	0.5-0.6	4
Crackers, unspecified type ⁴⁹	1.7	NA	NA
Crackers, wheat, shredded ⁴⁷	1.2	NA	2
Crackers, wheat, thin ⁴⁷	0.4	NA	2
Grains, cooked			
Barley, native ^{10,50,74,84,86}	3.4±0.8	1.7-4.2	24
Barley, high amylose ⁸⁶	3.5±0.3	3.1-3.8	18
Barley, low amylose, waxy ⁸⁶	0.3±0.0	0.3-0.3	18
Corn, hominy grits ⁸²	1.3	NA	NA
Corn, meal ^{76,79,82}	0.8±0.2	0.7-1.0	7
Corn, tamale ⁸⁷	1.3±0.4	0.9-1.6	6
Groats, buckwheat ^{88,89}	0.9±0.2	0.8-1.8	25
Millet ^{10,79}	1.0±0.5	0.8-1.6	4
Legumes			
Cooked			
Butter or lima bean ^{15,50}	6.4±1.8	1.2-7.1	10
Black bean ^{69,90-93}	2.7±1.7	1.2-5.3	12
Chickpea ^{45,50,52,76,90,93-99}	2.1±1.0	0.8-4.1	67
Fava bean ⁹²	0.7	NA	NA
Great Northern bean ^{92,93}	1.2±0.4	0.8-1.6	3
Kidney bean ^{15,50,51,79,93,100}	3.8±1.4	0.8-4.7	50
Kidney, white bean ^{50,92}	3.6±3.2	1.4-8.3	4
Lentils ^{10,50,52,76,79,94,95,98,101-104}	2.0±16.9	0.5-3.2	232
Lentil, gram dhal ¹⁰⁵	1.6±0.4	1.2-2.2	4
Mung bean ^{45,99,106}	1.2±0.3	0.6-1.5	20
Navy bean ^{92,95}	1.6±0.2	1.6-1.9	4
Peas ^{10,50,75,76,79,95,98,100,103}	1.9±0.9	0.9-6.3	40
Pinto or common brown bean ^{50,52,91,93,95,97,98,107}	2.0±0.3	1.4-2.4	27
Pinto, refried bean ⁹⁰	0.8	NA	4
Red, small bean ⁹³	1.4	NA	NA
White bean ^{52,84,93,103,108}	2.2±1.5	1.4-2.6	24

(continued on next page)

Table. Naturally occurring resistant starch in whole foods commonly consumed in the United States, extracted from peer-reviewed articles published from 1982 to September 2018 (*continued*)

Food	g RS ^a per 100 g Food ^b		
	Weighted mean±weighted SD ^{cd}	Range ^e	n ^f
Cooked and stored for 24 to 96 h			
Black ^{69,91}	4.7±2.0	2.3-7.3	21
Pinto or common brown ^{52,91,107}	2.7±1.6	2.0-8.4	43
Cooked, chilled, and reheated			
Lentils ^{102,109}	1.6±0.2	1.5-3.0	198
Noodles and pasta			
Noodles, egg, cooked ⁴⁷	0.2±0.2	0.0-0.4	4
Noodles, rice, cooked ⁴⁷	0.6±0.1	0.5-0.6	4
Pasta, corn, cooked ⁴¹	0.1	NA	8
Pasta, cooked ^{10,14,47,50,52,74,76,81,82}	1.5±0.8	0.2-2.9	34
Pasta, cooked then chilled ^{10,76}	1.0±0.2	0.8-1.4	5
Pasta, durum wheat, cooked ^{41,78,110-112}	1.2±1.2	0.2-3.2	23
Pasta, durum wheat, cooked then chilled 3 to 5 d ¹¹¹	3.4±0.0	3.4-3.5	12
Pasta, mixed grain, cooked ⁴¹	0.5	NA	6
Pasta, whole wheat, cooked ⁴⁷	0.2	NA	2
Other			
Hazelnuts ¹¹³	1.4±0.1	1.3-1.4	6
Potatoesⁱ			
Boiled, baked, or microwaved			
Potato, unknown variety ^{10,13,14,41,45,50,52,76,80,81,114-119}	0.7±0.5	0.1-2.0	74
Potato, high-amylose variety ¹¹⁸	4.6	NA	3
Potato, red variety ^{120,121}	1.7±1.3	0.7-3.8	16
Potato, russet ¹²¹	3.1±0.5	2.6-3.5	6
Potato, sweet variety ^{45,50,80}	0.5±0.5	0.3-2.1	15
Potato, yellow variety ^{120,121}	1.4±1.3	0.3-3.5	16
Cooked then chilled			
Potato, unknown variety ^{10,13,14,80,114-116,118,119}	1.3±0.6	0.3-2.4	47
Potato, high-amylose variety ¹¹⁸	6.4±0.2	6.0-6.8	6
Potato, red variety ^{120,121}	2.0±1.7	1.1-4.8	13
Potato, russet ¹²¹	4.3±0.5	3.8-4.7	6
Potato, salad ^{81,115}	5.2±1.9	1.0-5.9	7
Potato, sweet variety ⁸⁰	0.4±0.0	0.4-0.4	2
Potato, yellow variety ^{120,121}	2.5±3.8	0.6-5.4	16
Cooked then frozen			
Potato, unknown variety ¹¹⁷	0.4±0.0	0.4-0.4	2
Cooked, chilled then reheated			
Potato, unknown variety ^{115,116}	1.6±0.7	0.4-2.2	11
Potato, red variety ¹²¹	3.2±0.7	2.6-3.8	6
Potato, russet ¹²¹	3.9±0.3	3.6-4.2	6
Potato, yellow variety ¹²¹	5.1±2.3	2.2-4.3	6

(continued on next page)

Table. Naturally occurring resistant starch in whole foods commonly consumed in the United States, extracted from peer-reviewed articles published from 1982 to September 2018 (*continued*)

Food	g RS ^a per 100 g Food ^b		n ^f
	Weighted mean±weighted SD ^{cd}	Range ^e	
Fried			
Potato, unknown variety ^{80,81,114,115}	3.8±1.9	0.2-5.5	15
Potato, sweet variety ⁸⁰	0.3±0.0	0.3-0.3	2
Fried then frozen			
Potato, unknown variety ¹²²	2.9±0.6	2.3-3.8	12
Other			
Potato, chips ^{50,52,114,115}	3.1±0.7	1.4-4.5	11
Potato, instant ^{10,50,81,114,115}	1.4±1.1	0.2-2.4	13
Rice			
Steamed, boiled, or pressure cooked			
Brown ⁵⁰	0.1±0.0	0.0-0.1	5
White, long grain ^{5,14,50,52,81,84,123-125}	1.4±1.0	0.0-3.7	46
White, medium grain ^{5,41}	0.2±0.1	0.1-0.5	18
White, short grain ^{5,123,124}	0.3±0.2	0.0-0.6	20
White, high amylose ^{126,127}	0.8±0.1	0.7-0.9	18
White, intermediate amylose ^{126,127}	0.5±0.0	0.5-0.6	9
White, low amylose ^{124,126,127}	0.2±0.1	0.1-0.4	11
Cooked then chilled			
White, long grain ^{5,125}	0.9±0.8	0.4-2.6	13
White, medium grain ⁵	0.6±0.3	0.3-0.9	6
White, short grain ⁵	0.4±0.3	0.2-0.8	6
Stir-fried			
White, long grain ¹²⁴	2.2	—	2
White, short grain ¹²⁴	1.4±0.1	1.4-1.4	4
Stir-fried then chilled			
White, long grain ¹²⁴	4.5±0.8	3.9-5.2	4
White, short grain ¹²⁴	3.7±1.0	2.5-5.0	8

^aRS=resistant starch.

^bRS value in whole food with moisture included. For references providing RS values on a dry weight basis, the formula (1-% moisture)*% dry weight RS value was used to calculate RS in the whole food. Moisture content was either measured and provided in the reference or obtained from the US Department of Agriculture FoodData Central database (www.fdc.nal.usda.gov)⁴⁰ using the most similar food item.

^cSD=standard deviation.

^dFirst, the weighted mean RS was calculated for each food item by multiplying the mean RS in the whole food (with moisture) by the number of samples analyzed in the reference. Next, the individual weighted means were summed and divided by the total of samples from all references in the food category to obtain the overall weighted mean. Then, the weighted SD was calculated by the individual food item by multiplying the sample size by the mean RS of each whole food (with moisture) minus the overall weighted mean squared. Next, the individual weighted SD were summed and divided by the total sample size minus 1. Lastly, the square root of this value provided the overall weighted SD. The weighted SD is not provided if the SD was not reported in the reference, even if more than one sample was analyzed.

^eMinimum to maximum weighted mean RS values in each food item category with more than 1 mean RS value.

^fTotal number of samples analyzed for each food item.

^gNA=not available.

^hBaked corn tortilla moisture content was not available in the FoodData Central database,⁴⁰ therefore, the measured moisture from Rohlfing and colleagues⁶³ was used.

ⁱAnalyzed without skin.

stored at 4°C for 14 (4.7 g/100 g) and 7 (4.6 g/100 g) days, respectively; and cooked high amylose potatoes (4.6 g/100 g). The foods with the lowest mean RS content include multi-grain flaked store-bought cereal (0.0 g/100 g), cooked

bananas (0.0 g/100 g), and cooked unripe plantain crisps (0.0 g/100 g), fruit scones (0.1 g/100 g), wheat-germ bread (0.1 g/100 g), store-bought granola (0.1 g/100 g), cooked corn pasta (0.1 g/100 g), and cooked brown rice (0.1 g/100 g).

DISCUSSION

RS is found in many starch-containing foods, such as breads, grains, pasta, cereals, beans and legumes, potatoes, and rice. Over the last several years, an increase in the quantitative analysis of RS in foods resulted from the development of a commercially available assay that indirectly measures RS. A complete database showing the RS content of whole foods has not been developed because the commercially available assay was used widely by the scientific community. In addition, the prior reference providing RS content in foods included RS values on a dry weight basis,³⁹ which would not be accurate in whole foods, where moisture is included. Therefore, a database was developed based on data from peer-reviewed publications that analyzed the RS content in whole foods that are consumed by individuals in the United States.

The database has the potential to assist registered dietitian nutritionists, researchers, the food industry, and consumers in several ways. First, the database could be a resource for registered dietitian nutritionists to assist client and patient efforts to incorporate naturally occurring RS foods into their diets. Next, the database could be used as a resource for practitioners and researchers to support the development of well-controlled dietary intervention studies that may examine the appropriate amount of naturally derived RS from foods needed to achieve a specific health outcome. Studies have indicated that adequate amounts of RS (approximately ≥ 15 g/day)¹²⁸ can improve postprandial glucose^{26,36} and insulin sensitivity, especially among adults with insulin resistance,^{29,129} and modulate the gut microbiome,¹³⁰ which may improve gastrointestinal health. However, many of these studies used RS as a functional ingredient either alone or added to a specific food (eg, muffin or cookie) instead of foods with naturally occurring RS. Next, food-manufacturing companies can utilize the database to formulate new products that contain naturally occurring RS-containing foods. Lastly, the database can also be used as a resource for consumers who are interested in adding RS foods to their diet.

Variation in RS Content in Foods

The variation in RS content within and between each food item category can be influenced by many factors. Breeding techniques can increase the amylose to amylopectin ratio of the starch granule to allow for RS formation during the cooling process.⁴ According to the database, high-amylose grains have more than 10 times, boiled high-amylose potatoes have more than 4 times, and high-amylose cooked white rice has more than 4 times the amount of RS than low-amylose varieties.

Varietal differences among the same food also contain different amounts of RS.⁴ For example, according to the database, russet potatoes have a higher mean RS (4.3 g/100 g) than yellow (2.5 g/100 g) and red (2.0 g/100 g) potatoes that were cooked and then chilled. Butter or lima beans (6.4 g/100 g), kidney beans (3.8 g/100 g), and white kidney beans (3.6 g/100 g) had the highest mean amount of RS, while fava beans (0.7 g/100 g), refried pinto beans (0.7 g/100 g), Great Northern beans (1.2 g/100 g), and mung beans (1.2 g/100 g) had the lowest amount of mean RS among the cooked legumes.

The preparation methods and storage conditions can change the RS content of food. Heat and moisture improve the gelatinization properties of the starch granule, which

upon cooling allows the linear amylose chains to pack tightly and resist digestive enzymes, a process called retrogradation.⁹ Cooking and then chilling of bananas and plantains, pasta, stir-fried rice, black and pinto beans, and potatoes increase the RS content. However, reheating after chilling reduces the RS content in bananas and plantains, legumes, and potatoes. In fact, cooking bananas eliminates all RS. Storage temperature and duration also influence RS formation in some foods, such as corn tortillas. Freshly baked corn tortillas have the lowest RS content compared to corn tortillas stored at 4°C, in which the RS content increases with time. The maximum RS formation in corn tortillas occurs when stored at 4°C for 3 to 5 days. Another example includes durum wheat pasta, where chilling the cooked pasta for 3 to 5 days increases the RS by almost three times compared to the freshly cooked pasta. Baking temperature and duration of the baking process also influence RS formation in breads. White wheat bread baked at a lower temperature (120°C) for longer duration (20 hours) contained 2.2 g/100 g RS compared to 1.1 g/100 g RS when bread is baked a higher baking temperature (150°C) for shorter duration (3 hours) (specific data not shown in the Table).^{61,80}

Finally, the type of analytical methods used to quantify the amount of RS can vary across the same food. The differences in analytical methods used pose challenges in adequately quantifying RS, especially when RS is to be included as a component of dietary fiber declarations. For example, AOAC methods 2011.25 and 991.43 yield distinctly different fiber values for some RS sources.¹³¹ These analytical challenges can create confusion for consumers seeking to add sources of RS to their diet, especially when RS is incorporated into the dietary fiber quantity on the Nutrition Facts label.

RS as Functional Ingredient

Enrichment of processed foods with RS can improve RS intake beyond the consumption of naturally occurring RS foods. Adding RS as a functional ingredient, specifically RS2 and RS3, to foods can improve the nutritional profile and retain sensory appeal without significant changes in texture or shelf life.¹³²⁻¹³⁵ Apart from being bland in flavor and having a neutral white color, commercial sources of RS2 and RS3 have a high gelatinization temperature, finer particle size, and are less susceptible to changes during processing and storage than RS found naturally in foods.¹³⁵ These properties can improve texture, provide better mouthfeel, color, and food flavor, thereby increasing the consumer acceptability of foods with added RS.¹³⁵ For these reasons, enriching foods with RS2 or RS3 as a functional ingredient may be more advantageous than foods with naturally occurring RS because the commercial RS retains resistance under processing and cooking conditions.¹³⁶ For example, heating potatoes in water increases the solubility of the RS2 starch granule to improve digestibility and reduce resistance, thus lowering RS content. However, RS as a functional ingredient incorporated into pasta will retain its resistance when cooked in water.

RS Food Claims Based on Clinical Trials

The US Food and Drug Administration (FDA) has now recognized some types of RS, specifically high amylose starch containing RS2, as dietary fibers for Nutrition Facts and

Supplement Facts labeling.¹³⁷ In addition, the US FDA authorized a qualified health claim for high-amylose maize RS and reduction of risk of type 2 diabetes. Additional requirements for foods bearing the claim are described in FDA Letter of Enforcement Discretion: Docket # FDA-2015-Q-2352.¹³⁸ The FDA will exercise enforcement discretion for the following qualified health claim statements: “High-amylose maize resistant starch may reduce the risk of Type 2 diabetes. FDA has concluded that there is limited scientific evidence for this claim.”¹³⁸ “High-amylose maize resistant starch, a type of fiber, may reduce the risk of Type 2 diabetes. FDA has concluded that there is limited scientific evidence for this claim.”¹³⁸

The clinical science for this qualified health claim is based on improved insulin sensitivity in healthy or at-risk individuals after consuming high-amylose maize RS.^{29,30,37,129,139,140} The European Food Safety Authority has also authorized a health claim that describes the benefit of RS on postprandial blood glucose concentrations.¹³⁷

Several strengths and limitations of this review must be mentioned. The strengths include identifying RS in foods that were analyzed in samples appropriate for human consumption. For example, raw foods (except bananas, plantains, and oats) or starches were not included, as they are not suitable for consumption, although some individuals add RS flour to foods as a functional ingredient. However, this review did not include these type of flours because they are manufactured and not present naturally in foods. In addition, three databases were searched by hand to include as many peer-reviewed articles quantifying RS in foods as possible. The database adjusted the RS value to include moisture if given as percent dry weight basis, which is more representative of RS in whole foods. A weighted sampling method was used to provide more weight for values with higher sampling representation compared to lower, which improves the validity of the data.

A few limitations must also be noted. The method of RS analysis highlights one major limitation. There is an ever-present gap between *in vitro* and *in vivo* analysis regardless of the methods used to quantify RS. Englyst and colleagues,^{7,8} and many others who adapted the Englyst procedure, based their RS analysis on a broad assumption of homogenization or milling food showing similar properties as those of chewing. However, chewing is very individualized and cannot be controlled to replicate a milled or homogenized product. Estimation of newly identified RS, such as RS4, deviates from methods utilized to estimate RS2 and RS3. The changes range from significant decrease in initial incubation times to a cold KOH treatment in order to solubilize RS4.¹⁴¹ It is clear that RS estimation is a constantly evolving process and any comparisons of RS values by different analytical methods are bound to reflect some amount of over- or underestimation. Also, a method to quantify RS5 has not been described. Another limitation was the adjustment of moisture when RS was provided on a percent dry weight basis. If the moisture was not measured and given in the reference, a food database was utilized to identify the moisture content in a similar food item. In addition, environmental and natural genetic differences influence starch digestibility and are inherently difficult to control.⁴ Processing, cooking, and storing methods can change the properties of the starch granule to allow for more or less resistance. Future research attempting to standardize RS quantification is needed to be

more accurate in assessing RS content in foods. Lastly, only peer-reviewed articles estimating RS in foods were included. Private or food companies conducting an independent RS analysis in foods were excluded, which limited the comprehensiveness of the database.

CONCLUSIONS

RS is found naturally in both processed and whole starchy foods, including breads, cereals, bananas and plantains, grains, noodles and pasta, potatoes, rice, and legumes. According to the database, raw foods, including oats and plantains, had the highest RS content. Among cooked foods, potatoes and grains (barley and rice) bred to have a higher amylose to amylopectin ratio have higher amounts of RS than those with a lower amylose to amylopectin ratio. Potatoes and grains that are cooked and then chilled have more RS than if boiled or heated, where the chilling process promotes the retrogradation of the starch granule to make it less digestible. The duration of storage also increases RS in some foods, such as corn tortillas, durum wheat pasta, and black and pinto beans. Foods with the lowest RS include multigrain flaked cereal, cooked bananas, cooked unripe plantain chips, fruit scones, wheat-germ bread, store-bought granola, cooked corn pasta, and cooked brown rice.

References

- Dahl WJ, Stewart ML. Position of the Academy of Nutrition and Dietetics: Health implications of dietary fiber. *J Acad Nutr Diet.* 2015;115(11):1861-1870.
- Grooms KN, Ommerborn MJ, Pham DQ, Djousse L, Clark CR. Dietary fiber intake and cardiometabolic risks among US adults, NHANES 1999-2010. *Am J Med.* 2013;126(12):1059-1067. e1051-1054.
- Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids.* Washington, DC: National Academies Press; 2005.
- Birt DF, Boylston T, Hendrich S, et al. Resistant starch: Promise for improving human health. *Adv Nutr.* 2013;4(6):587-601.
- Chiu YT, Stewart ML. Effect of variety and cooking method on resistant starch content of white rice and subsequent postprandial glucose response and appetite in humans. *Asia Pac J Clin Nutr.* 2013;22(3):372-379.
- Westenbrink S, Brunt K, van der Kamp JW. Dietary fibre: Challenges in production and use of food composition data. *Food Chem.* 2013;140(3):562-567.
- Englyst H, Wiggins HS, Cummings JH. Determination of the non-starch polysaccharides in plant foods by gas-liquid chromatography of constituent sugars as alditol acetates. *Analyst.* 1982;107(1272):307-318.
- Englyst HN, Cummings JH. Digestion of the polysaccharides of some cereal foods in the human small intestine. *Am J Clin Nutr.* 1985;42(5):778-787.
- Perera A, Meda V, Tyler RT. Resistant starch: A review of analytical protocols for determining resistant starch and of factors affecting the resistant starch content of foods. *Food Res Int.* 2010;43(8):1959-1974.
- Englyst HN, Kingman SM, Cummings JH. Classification and measurement of nutritionally important starch fractions. *Eur J Clin Nutr.* 1992;46(suppl 2):S33-S50.
- Berry CS. Resistant starch: Formation and measurement of starch that survives exhaustive digestion with amylolytic enzymes during the determination of dietary fibre. *J Cereal Sci.* 1986;4(4):301-314.
- Goñi I, García-Diz L, Mañas E, Saura-Calixto F. Analysis of resistant starch: A method for foods and food products. *Food Chem.* 1996;56(4):445-449.
- Muir JG, O’Dea K. Measurement of resistant starch: Factors affecting the amount of starch escaping digestion *in vitro*. *Am J Clin Nutr.* 1992;56(1):123-127.

14. Akerberg AK, Liljeberg HG, Granfeldt YE, Drews AW, Björck IM. An in vitro method, based on chewing, to predict resistant starch content in foods allows parallel determination of potentially available starch and dietary fiber. *J Nutr*. 1998;128(3):651-660.
15. McCleary BV, Monaghan DA. Measurement of resistant starch. *J AOAC Int*. 2002;85(3):665-675.
16. Megazyme. Resistant starch assay kit. www.megazyme.com. Published 2019. Accessed June 16, 2019.
17. Shen D, Bai H, Li Z, Yu Y, Zhang H, Chen L. Positive effects of resistant starch supplementation on bowel function in healthy adults: A systematic review and meta-analysis of randomized controlled trials. *Int J Food Sci Nutr*. 2017;68(2):149-157.
18. Grabitske HA, Slavin JL. Low-digestible carbohydrates in practice. *J Am Diet Assoc*. 2008;108(10):1677-1681.
19. Purwani EY, Purwadaria T, Suhartono MT. Fermentation RS3 derived from sago and rice starch with *Clostridium butyricum* BCC B2571 or *Eubacterium rectale* DSM 17629. *Anaerobe*. 2012;18(1):55-61.
20. Zaman SA, Sarbini SR. The potential of resistant starch as a prebiotic. *Crit Rev Biotechnol*. 2016;36(3):578-584.
21. Martinez I, Kim J, Duffy PR, Schlegel VL, Walter J. Resistant starches types 2 and 4 have differential effects on the composition of the fecal microbiota in human subjects. *PLoS One*. 2010;5(11):e15046.
22. Alfa MJ, Strang D, Tappia PS, et al. A randomized trial to determine the impact of a digestion resistant starch composition on the gut microbiome in older and mid-age adults. *Clin Nutr*. 2018;37(3):797-807.
23. Maier TV, Lucio M, Lee LH, et al. Impact of dietary resistant starch on the human gut microbiome, metaproteome, and metabolome. *MBio*. 2017;8(5).
24. Willis HJ, Eldridge AL, Beiseigel J, Thomas W, Slavin JL. Greater satiety response with resistant starch and corn bran in human subjects. *Nutr Res*. 2009;29(2):100-105.
25. Ble-Castillo JL, Juarez-Rojop IE, Tovilla-Zarate CA, et al. Acute consumption of resistant starch reduces food intake but has no effect on appetite ratings in healthy subjects. *Nutrients*. 2017;9(7).
26. Maziarz MP, Preisendanz S, Juma S, Imrhan V, Prasad C, Vijayagopal P. Resistant starch lowers postprandial glucose and leptin in overweight adults consuming a moderate-to-high-fat diet: A randomized-controlled trial. *Nutr J*. 2017;16(1):14.
27. Zenel AM, Stewart ML. High amylose white rice reduces postprandial glycemic response but not appetite in humans. *Nutrients*. 2015;7(7):5362-5374.
28. Johnston KL, Thomas EL, Bell JD, Frost GS, Robertson MD. Resistant starch improves insulin sensitivity in metabolic syndrome. *Diabet Med*. 2010;27(4):391-397.
29. Maki KC, Pelkman CL, Finocchiaro ET, et al. Resistant starch from high-amylose maize increases insulin sensitivity in overweight and obese men. *J Nutr*. 2012;142(4):717-723.
30. Robertson MD, Bickerton AS, Dennis AL, Vidal H, Frayn KN. Insulin-sensitizing effects of dietary resistant starch and effects on skeletal muscle and adipose tissue metabolism. *Am J Clin Nutr*. 2005;82(3):559-567.
31. Peterson CM, Beyl RA, Marlatt KL, et al. Effect of 12 wk of resistant starch supplementation on cardiometabolic risk factors in adults with prediabetes: A randomized controlled trial. *Am J Clin Nutr*. 2018.
32. Wutzke KD, Schmidek KV. The effect of resistant starches on fat oxidation in healthy adults as measured by a (13)CO2-breath test. *Isotopes Environ Health Stud*. 2017;53(6):553-562.
33. Yuan HC, Meng Y, Bai H, Shen DQ, Wan BC, Chen LY. Meta-analysis indicates that resistant starch lowers serum total cholesterol and low-density cholesterol. *Nutr Res*. 2018;54:1-11.
34. Lockyer S, Nugent AP. Health effects of resistant starch. *Nutr Bull*. 2017;42(1):10-41.
35. Bergeron N, Williams PT, Lamendella R, et al. Diets high in resistant starch increase plasma levels of trimethylamine-N-oxide, a gut microbiome metabolite associated with CVD risk. *Br J Nutr*. 2016;116(12):2020-2029.
36. Stewart ML, Zimmer JP. A high fiber cookie made with resistant starch type 4 reduces post-prandial glucose and insulin responses in healthy adults. *Nutrients*. 2017;9(3).
37. Gower BA, Bergman R, Stefanovski D, et al. Baseline insulin sensitivity affects response to high-amylose maize resistant starch in women: A randomized, controlled trial. *Nutr Metab (Lond)*. 2016;13:2.
38. O'Keefe SJ, Chung D, Mahmoud N, et al. Why do African Americans get more colon cancer than Native Africans? *J Nutr*. 2007;137(1 suppl):175S-182S.
39. Murphy MM, Douglass JS, Birkett A. Resistant starch intakes in the United States. *J Am Diet Assoc*. 2008;108(1):67-78.
40. US Department of Agriculture ARS. FoodData Central. www.fdc.nal.usda.gov. Published 2019. Accessed October 1, 2019.
41. Carcea M, Salvatorelli S, Turfani V. Measurement of resistant starch in cooked cereal-based foods. *Qual Assur Saf Crop*. 2009;1(4):240-245.
42. Odenigbo AM, Asumugha VU, Ubbor S, Ngadi M. In vitro starch digestibility of plantain and cooking-banana at ripe and unripe stages. *Int Food Res J*. 2013;20(6):3027-3031.
43. Lintas C, Cappelloni M, Adorisio S, Clementi A, Del Toma E. Effect of ripening on resistant starch and total sugars in *Musa paradisiaca sapientum*: Glycaemic and insulinaemic responses in normal subjects and NIDDM patients. *Eur J Clin Nutr*. 1995;49(suppl 3):S303-S306.
44. Englyst HN, Cummings JH. Digestion of the carbohydrates of banana (*Musa paradisiaca sapientum*) in the human small intestine. *Am J Clin Nutr*. 1986;44(1):42-50.
45. Patel K, Shurpalekar KS. Resistant starch content of Indian foods. *Plant Food Hum Nutr*. 1994;45:91-95.
46. Oladele EO, Williamson G. Impact of resistant starch in three plantain (*Musa AAB*) products on glycaemic response of healthy volunteers. *Eur J Nutr*. 2016;55(1):75-81.
47. Gelroth JA, Ranhotra GS, Gelroth JA, Ranhotra GS. Determination of resistant starch in selected grain-based foods. *J AOAC Int*. 2000;83(4):988-991.
48. Djurle S, Andersson AAM, Andersson R. Effects of baking on dietary fibre, with emphasis on β -glucan and resistant starch, in barley breads. *J Cereal Sci*. 2018;79:449-455.
49. Englyst KN, Vinoy S, Englyst HN, Lang V. Glycaemic index of cereal products explained by their content of rapidly and slowly available glucose. *Br J Nutr*. 2003;89(3):329-340.
50. Englyst HN, Veenstra J, Hudson GJ. Measurement of rapidly available glucose (RAG) in plant foods: A potential in vitro predictor of the glycaemic response. *Br J Nutr*. 1996;75(3):327-337.
51. Abdel-Aal E-SM, Rabalski I. Effect of baking on nutritional properties of starch in organic spelt whole grain products. *Food Chem*. 2008;111(1):150-156.
52. Saura-Calixto F, Garcia-Alonso A, Goni I, Bravo L. In vitro determination of the indigestible fraction in foods: An alternative to dietary fiber analysis. *J Agric Food Chem*. 2000;48(8):3342-3347.
53. Food Standards Agency. *Nutrient Analysis of Bread and Morning Goods. Project Number AN1062: Analytical Report*. London, UK: Ministry of Agriculture Fisheries and Food, Department of Health and the Scottish Executive; 2001.
54. Muir JG, Yeow EG, Keogh J, et al. Combining wheat bran with resistant starch has more beneficial effects on fecal indexes than does wheat bran alone. *Am J Clin Nutr*. 2004;79(6):1020-1028.
55. Giuberti G, Fortunati P, Gallo A. Can different types of resistant starch influence the in vitro starch digestion of gluten free breads? *J Cereal Sci*. 2016;70:253-255.
56. De La Hera E, Rosell CM, Gomez M. Effect of water content and flour particle size on gluten-free bread quality and digestibility. *Food Chem*. 2014;151:526-531.
57. Buddrick O, Jones OAH, Hughes JG, Kong I, Small DM. The effect of fermentation and addition of vegetable oil on resistant starch formation in wholegrain breads. *Food Chem*. 2015;180:181-185.
58. Collar C, Jimenez T, Conte P, Fadda C. Impact of ancient cereals, pseudocereals and legumes on starch hydrolysis and antiradical activity of technologically viable blended breads. *Carbohydr Polym*. 2014;113:149-158.
59. Ragaei S, Guzar I, Dhull N, Seetharaman K. Effects of fiber addition on antioxidant capacity and nutritional quality of wheat bread. *LWT Food Sci Technol*. 2011;44(10):2147-2153.
60. Giacco R, Brighenti F, Parillo M, et al. Characteristics of some wheat-based foods of the Italian diet in relation to their influence on

- postprandial glucose metabolism in patients with type 2 diabetes. *Br J Nutr*. 2001;85(1):33-40.
61. Amaral O, Guerreiro CS, Gomes A, Cravo M. Resistant starch production in wheat bread: Effect of ingredients, baking conditions and storage. *Eur Food Res Technol*. 2016;242(10):1747-1753.
 62. Rizzello CG, Calasso M, Campanella D, De Angelis M, Gobbetti M. Use of sourdough fermentation and mixture of wheat, chickpea, lentil and bean flours for enhancing the nutritional, texture and sensory characteristics of white bread. *Int J Food Microbiol*. 2014;180:78-87.
 63. Rohlffing KA, Paez A, Kim HJ, White PJ. Effects of resistant starch and fiber from high-amylose non-floury corn on tortilla texture. *Cereal Chem*. 2010;87(6):581-585.
 64. Bello-Perez LA, Flores-Silva PC, Agama-Acevedo E, de Dios Figueroa-Cardenas J, Lopez-Valenzuela JA, Campanella OH. Effect of the nixtamalization with calcium carbonate on the indigestible carbohydrate content and starch digestibility of corn tortilla. *J Cereal Sci*. 2014;60(2):421-425.
 65. Osorio-Díaz P, Agama-Acevedo E, Bello-Pérez LA, Islas-Hernández JJ, Gomez-Montiel NO, Paredes-López O. Effect of endosperm type on texture and in vitro starch digestibility of maize tortillas. *LWT Food Sci Technol*. 2011;44(3):611-615.
 66. Agama-Acevedo E, Rendon-Villalobos R, Tovar J, Paredes-Lopez O, Islas-Hernandez JJ, Bello-Perez LA. In vitro starch digestibility changes during storage of maize flour tortillas. *Die Nahrung*. 2004;48(1):38-42.
 67. Islas-Hernández JJ, Rendón-Villalobos R, Agama-Acevedo E, et al. In vitro digestion rate and resistant starch content of tortillas stored at two different temperatures. *LWT Food Sci Technol*. 2006;39(8):947-951.
 68. Rendon R, Arturo Bello-Pérez L, Osorio-Díaz P, Tovar J, Paredes-Lopez O. Effect of storage time on in vitro digestibility and resistant starch content of nixtamal, masa, and tortilla. *J Agric Food Chem*. 2005;53(4):1281-1285.
 69. Sayago-Ayerdi SG, Tovar J, Osorio-Díaz P, Paredes-Lopez O, Bello-Perez LA. In vitro starch digestibility and predicted glycemic index of corn tortilla, black beans, and tortilla-bean mixture: Effect of cold storage. *J Agric Food Chem*. 2005;53(4):1281-1285.
 70. Gutiérrez-Dorado R, Ayala-Rodríguez AE, Milán-Carrillo J, et al. Technological and nutritional properties of flours and tortillas from nixtamalized and extruded quality protein maize (*Zea mays L.*). *Cereal Chem*. 2008;85(6):808-816.
 71. Santiago-Ramos D, De Dios Figueroa-Cárdenas J, Véles-Medina JJ, et al. Resistant starch formation in tortillas from an ecological nixtamalization process. *Cereal Chem*. 2015;92(2):185-192.
 72. Angioloni A, Collar C. Physicochemical and nutritional properties of reduced-caloric density high-fibre breads. *LWT Food Sci Technol*. 2011;44(3):747-758.
 73. Skrabanja V, Liljeberg HG, Hedley CL, Kreft I, Bjorck IM. Influence of genotype and processing on the in vitro rate of starch hydrolysis and resistant starch formation in peas (*Pisum sativum L.*). *J Agric Food Chem*. 1999;47(5):2033-2039.
 74. Englyst KN, Englyst HN, Hudson GJ, Cole TJ, Cummings JH. Rapidly available glucose in foods: An in vitro measurement that reflects the glycemic response. *Am J Clin Nutr*. 1999;69(3):448-454.
 75. Marlatt KL, White UA, Beyl RA, et al. Role of resistant starch on diabetes risk factors in people with prediabetes: Design, conduct, and baseline results of the STARCH trial. *Contemp Clin Trials*. 2018;65:99-108.
 76. Rosin PM, Lajolo FM, Menezes EW. Measurement and characterization of dietary starches. *J Food Comp Anal*. 2002;15:367-377.
 77. Hallstrom E, Sestili F, Lafandra D, Bjorck I, Ostman E. A novel wheat variety with elevated content of amylose increases resistant starch formation and may beneficially influence glycaemia in healthy subjects. *Food Nutr Res*. 2011;55.
 78. Hoebler C, Karinthi A, Chiron H, Champ M, Barry JL. Bioavailability of starch in bread rich in amylose: Metabolic responses in healthy subjects and starch structure. *Eur J Clin Nutr*. 1999;53(5):360-366.
 79. Dodevska MS, Djordjevic BI, Sobajic SS, Miletic ID, Djordjevic PB, Dimitrijevic-Sreckovic VS. Characterisation of dietary fibre components in cereals and legumes used in Serbian diet. *Food Chem*. 2013;141(3):1624-1629.
 80. Yadav BS. Effect of frying, baking and storage conditions on resistant starch content of foods. *Br Food J*. 2011;113(6):710-719.
 81. Liljeberg Elmstahl H. Resistant starch content in a selection of starchy foods on the Swedish market. *Eur J Clin Nutr*. 2002;56(6):500-505.
 82. Bednar GE, Patil AR, Murray SM, Grieshop CM, Merchen NR, Fahey GC Jr. Starch and fiber fractions in selected food and feed ingredients affect their small intestinal digestibility and fermentability and their large bowel fermentability in vitro in a canine model. *J Nutr*. 2001;131(2):276-286.
 83. Olesen M, Rumessen JJ, Gudmand-Hoyer E. The hydrogen breath test in resistant starch research. *Eur J Clin Nutr*. 1992;46(suppl 2):S133-S134.
 84. Muir JG, O'Dea K. Validation of an in vitro assay for predicting the amount of starch that escapes digestion in the small intestine of humans. *Am J Clin Nutr*. 1993;57(4):540-546.
 85. Marlett JA, Longacre MJ. Comparison of in vitro and in vivo measures of resistant starch in selected grain products. *Cereal Chem*. 1996;73(1):63-68.
 86. Emami S, Meda V, Pickard MD, Tyler RT. Impact of micronization on rapidly digestible, slowly digestible, and resistant starch concentrations in normal, high-amylose, and waxy barley. *J Agric Food Chem*. 2010;58(17):9793-9799.
 87. Mariscal-Moreno RM, de Dios Figueroa Cárdenas J, Santiago-Ramos D, Rayas-Duarte P, Velés-Medina JJ, Martínez-Flores HE. Nixtamalization process affects resistant starch formation and glycemic index of tamales. *J Food Sci*. 2017;82(5):1110-1115.
 88. Skrabanja V, Liljeberg Elmstahl HG, Kreft I, Bjorck IM. Nutritional properties of starch in buckwheat products: Studies in vitro and in vivo. *J Agric Food Chem*. 2001;49(1):490-496.
 89. Lu L, Murphy K, Baik B-K. Genotypic variation in nutritional composition of buckwheat groats and husks. *Cereal Chem*. 2013;90(2):132-137.
 90. Fabbri ADT, Schacht RW, Crosby GA. Evaluation of resistant starch content of cooked black beans, pinto beans, and chickpeas. *NFS J*. 2016;3:8-12.
 91. Osorio-Díaz P, Bello-Pérez LA, Sayago-Ayerdi SG, Benítez-Reyes MdP, Tovar J, Paredes-López O. Effect of processing and storage time on in vitro digestibility and resistant starch content of two bean (*Phaseolus vulgaris L.*) varieties. *J Sci Food Agric*. 2003;83(12):1283-1288.
 92. Pujolà M, Farreras A, Casañas F. Protein and starch content of raw, soaked and cooked beans (*Phaseolus vulgaris L.*). *Food Chem*. 2007;102(4):1034-1041.
 93. Wang N, Hatcher DW, Tyler RT, Toews R, Gawalko EJ. Effect of cooking on the composition of beans (*Phaseolus vulgaris L.*) and chickpeas (*Cicer arietinum L.*). *Food Res Int*. 2010;43(2):589-594.
 94. Aguilera Y, Esteban RM, Benitez V, Molla E, Martin-Cabrejas MA. Starch, functional properties, and microstructural characteristics in chickpea and lentil as affected by thermal processing. *J Agric Food Chem*. 2009;57(22):10682-10688.
 95. Brummer Y, Kaviani M, Tosh SM. Structural and functional characteristics of dietary fibre in beans, lentils, peas and chickpeas. *Food Res Int*. 2015;67:117-125.
 96. Hawkins A, Johnson SK. In vitro carbohydrate digestibility of whole-chickpea and chickpea bread products. *Int J Food Sci Nutr*. 2005;56(3):147-155.
 97. Marconi E, Ruggeri S, Cappelloni M, Leonardi D, Carnovale E. Physicochemical, nutritional, and microstructural characteristics of chickpeas (*Cicer arietinum L.*) and common beans (*Phaseolus vulgaris L.*) following microwave cooking. *J Agric Food Chem*. 2000;48(12):5986-5994.
 98. De Almeida Costa GE, Da Silva Queiroz-Monici K, Pissini Machado Reis SM, De Oliveira AC. Chemical composition, dietary fibre and resistant starch contents of raw and cooked pea, common bean, chickpea and lentil legumes. *Food Chem*. 2006;94(3):327-330.
 99. Veena A, Urooj A, Puttaraj S. Effect of processing on the composition of dietary fibre and starch in some legumes. *Die Nahrung*. 1995;39(2):132-138.
 100. Eyaru R, Shrestha AK, Arcot J. Effect of various processing techniques on digestibility of starch in red kidney bean (*Phaseolus*

- vulgaris*) and two varieties of peas (*Pisum sativum*). *Food Res Int*. 2009;42(8):956-962.
101. Araya H, Pak N, Vera G, Alvina M. Digestion rate of legume carbohydrates and glycemic index of legume-based meals. *Int J Food Sci Nutr*. 2003;54(2):119-126.
 102. Johnson CR, Thavarajah D, Thavarajah P, Payne S, Moore J, Ohm JB. Processing, cooking, and cooling affect prebiotic concentrations in lentil (*Lens culinaris Medikus*). *J Food Comp Anal*. 2015;38:106-111.
 103. Piecyk M, Wolosiak R, Druzynska B, Worobiej E. Chemical composition and starch digestibility in flours from Polish processed legume seeds. *Food Chem*. 2012;135(3):1057-1064.
 104. Wang N, Hatcher DW, Toews R, Gawalko EJ. Influence of cooking and dehulling on nutritional composition of several varieties of lentils (*Lens culinaris*). *LWT Food Sci Technol*. 2009;42(4):842-848.
 105. Nigudkar MR, Madan JG. Resistant starch content of traditional Indian legume preparations. *Curr Res Nutr Food Sci*. 2017;5(3).
 106. Kaur M, Sandhu KS, Ahlawat R, Sharma S. In vitro starch digestibility, pasting and textural properties of mung bean: Effect of different processing methods. *J Food Sci Technol*. 2015;52(3):1642-1648.
 107. Landa-Habana L, Pina-Hernandez A, Agama-Acevedo E, Tovar J, Bello-Perez LA. Effect of cooking procedures and storage on starch bioavailability in common beans (*Phaseolus vulgaris L.*). *Plant Foods Hum Nutr*. 2004;59(4):133-136.
 108. Noah L, Guillon F, Bouchet B, et al. Digestion of carbohydrate from white beans (*Phaseolus vulgaris L.*) in healthy humans. *J Nutr*. 1998;128(6):977-985.
 109. Siva N, Thavarajah P, Thavarajah D. The impact of processing and cooking on prebiotic carbohydrates in lentil. *J Food Comp Anal*. 2018;70:72-77.
 110. Aravind N, Sissons M, Fellows CM, Blazek J, Gilbert EP. Optimisation of resistant starch II and III levels in durum wheat pasta to reduce in vitro digestibility while maintaining processing and sensory characteristics. *Food Chem*. 2013;136(2):1100-1109.
 111. Bello-Perez LA, Flores-Silva PC, Camelo-Mendez GA, Paredes-Lopez O, Figueroa-Cardenas JD. Effect of the nixtamalization process on the dietary fiber content, starch digestibility, and antioxidant capacity of blue maize tortilla. *Cereal Chem*. 2015;92(3):265-270.
 112. Khan I, Yousif A, Johnson SK, Gamalath S. Effect of sorghum flour addition on resistant starch content, phenolic profile and antioxidant capacity of durum wheat pasta. *Food Res Int*. 2013;54(1):578-586.
 113. Kalkan F, Vanga SK, Garipey Y, Raghavan V. Effect of MW-assisted roasting on nutritional and chemical properties of hazelnuts. *Food Nutr Res*. 2015;59:28916.
 114. Garcia-Alonso A, Goni I. Effect of processing on potato starch: In vitro availability and glycaemic index. *Die Nahrung*. 2000;44(1):19-22.
 115. Kingman SM, Englyst HN. The influence of food preparation methods on the in-vitro digestibility of starch in potatoes. *Food Chem*. 1994;49(2):181-186.
 116. Leeman M, Ostman E, Björck I. Vinegar dressing and cold storage of potatoes lowers postprandial glycaemic and insulinaemic responses in healthy subjects. *Eur J Clin Nutr*. 2005;59(11):1266-1271.
 117. Englyst HN, Cummings JH. Digestion of polysaccharides of potato in the small intestine of man. *Am J Clin Nutr*. 1987;45(2):423-431.
 118. Zhao X, Andersson M, Andersson R. Resistant starch and other dietary fiber components in tubers from a high-amylose potato. *Food Chem*. 2018;251:58-63.
 119. Monro J, Mishra S, Blandford E, Anderson J, Genet R. Potato genotype differences in nutritionally distinct starch fractions after cooking, and cooking plus storing cool. *J Food Comp Anal*. 2009;22(6):539-545.
 120. Larder CE, Abergel M, Kubow S, Donnelly DJ. Freeze-drying affects the starch digestibility of cooked potato tubers. *Food Res Int*. 2018;103:208-214.
 121. Raatz SK, Idso L, Johnson LK, Jackson MI, Combs GF. Resistant starch analysis of commonly consumed potatoes: Content varies by cooking method and service temperature but not by variety. *Food Chem*. 2016;208:297-300.
 122. de Vasconcelos NCM, Salgado SM, Livera AVS, de Andrade SAC, de Oliveira MG, Stamford TLM. Influence of heat treatment on the sensory and physical characteristics and carbohydrate fractions of french-fried potatoes (*Solanum tuberosum L.*). *Food Sci Technol*. 2015;35(3):561-569.
 123. Ortuno J, Ros G, Periago MJ, Martinez C, Lopez G. Cooking water uptake and starch digestible value of selected spanish rices. *J Food Qual*. 1996;19(1):79-89.
 124. Reed MO, Ai Y, Leutcher JL, Jane JL. Effects of cooking methods and starch structures on starch hydrolysis rates of rice. *J Food Sci*. 2013;78(7):H1076-H1081.
 125. Sonia S, Witjaksono F, Ridwan R. Effect of cooling of cooked white rice on resistant starch content and glycemic response. *Asia Pac J Clin Nutr*. 2015;24(4):620-625.
 126. Chen MH, Bergman CJ, McClung AM, Everette JD, Tabien RE. Resistant starch: Variation among high amylose rice varieties and its relationship with apparent amylose content, pasting properties and cooking methods. *Food Chem*. 2017;234:180-189.
 127. Sagum R, Arcot J. Effect of domestic processing methods on the starch, non-starch polysaccharides and in vitro starch and protein digestibility of three varieties of rice with varying levels of amylose. *Food Chem*. 2000;70(1):107-111.
 128. Maziarz MP. Role of fructans and resistant starch in diabetes care. *Diabetes Spectrum*. 2013;26(1):35-39.
 129. Robertson MD, Wright JW, Loizon E, et al. Insulin-sensitizing effects on muscle and adipose tissue after dietary fiber intake in men and women with metabolic syndrome. *J Clin Endocrinol Metab*. 2012;97(9):3326-3332.
 130. Venkataraman A, Sieber JR, Schmidt AW, Waldron C, Theis KR, Schmidt TM. Variable responses of human microbiomes to dietary supplementation with resistant starch. *Microbiome*. 2016;4(1):33.
 131. McCleary BV, Sloane N, Draga A. Determination of total dietary fibre and available carbohydrates: A rapid integrated procedure that simulates in vivo digestion. *Starch Stärke*. 2015;67(9-10):860-883.
 132. Maziarz M, Sherrard M, Juma S, Prasad C, Imrhan V, Vijayagopal P. Sensory characteristics of high-amylose maize-resistant starch in three food products. *Food Sci Nutr*. 2013;1(2):117-124.
 133. Baixauli R, Salvador A, Martinez-Cervera S, Fiszman S. Distinctive sensory features introduced by resistant starch in baked products. *LWT Food Sci Technol*. 2008;41(10):1927-1933.
 134. Laguna L, Varela P, Salvador A, Sanz T, Fiszman SM. Balancing texture and other sensory features in reduced fat short-dough biscuits. *J Text Stud*. 2012;43(3):235-245.
 135. Fuentes-Zaragoza E, Riquelme-Navarrete M, Sánchez-Zapata E, Pérez-Álvarez J. Resistant starch as functional ingredient: A review. *Food Res Int*. 2010;43(4):931-942.
 136. Parada J, Aguilera JM. In vitro digestibility and glycemic response of potato starch is related to granule size and degree of gelatinization. *J Food Sci*. 2009;74(1):E34-E38.
 137. Scientific opinion on the substantiation of health claims related to resistant starch and reduction of post-prandial glycaemic responses (ID 681), "digestive health benefits" (ID 682) and "favours a normal colon metabolism" (ID 783) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA J*. 2011;9(4):2024.
 138. Food and Drug Administration, Department of Health and Human Services. *High-Amylose Starch and Diabetes*. Docket Number FDA-2015-Q-2352. College Park, MD: Food and Drug Administration; 2015.
 139. Bodinham CL, Frost GS, Robertson MD. Acute ingestion of resistant starch reduces food intake in healthy adults. *Br J Nutr*. 2010;103(6):917-922.
 140. Dainty SA, Klingel SL, Pilkey SE, et al. Resistant starch bagels reduce fasting and postprandial insulin in adults at risk of type 2 diabetes. *J Nutr*. 2016;146(11):2252-2259.
 141. Shi J, Sun Z, Shi YC. Improved in vitro assay of resistant starch in cross-linked phosphorylated starch. *Carbohydr Polym*. 2019;210:210-214.

AUTHOR INFORMATION

M. A. Patterson is an assistant professor and M. Maiya is a research coordinator for the Office of Research and Sponsored Programs, Department of Nutrition and Food Sciences, Texas Woman's University, Houston. M. L. Stewart is global research and development technical director of plant-based proteins, Ingredion Incorporated, Bridgewater, NJ.

Address correspondence to: Mindy A. Patterson, PhD, RDN, Department of Nutrition and Food Sciences, Texas Woman's University, 6700 Fannin St, Houston, TX 77030. E-mail: mpatterson14@twu.edu

STATEMENT OF POTENTIAL CONFLICT OF INTEREST

M. L. Stewart is an employee of Ingredion Incorporated; however, the perspectives provided by the author are her own, and do not necessarily reflect the perspectives of her employer. No potential conflict of interest was reported by the remaining authors.

FUNDING/SUPPORT

There is no funding to disclose.

AUTHOR CONTRIBUTIONS

M. A. Patterson developed the idea. M. A. Patterson and M. Maiya identified and extracted the peer-reviewed articles meeting inclusion criteria. M. A. Patterson and M. Maiya developed and contributed to the database. M. A. Patterson, M. Maiya, and M. L. Stewart wrote the initial manuscript, revised and commented on all versions, and approved the final version of the manuscript.